

AD-A061 321

CALIFORNIA UNIV LOS ANGELES DEPT OF CHEMISTRY

F/6 7/4

A B(11) NMR STUDY OF THE CLOSO-HYDRIDORHODACARBORANE 3,3-(C2H5--ETC(U)

JAN 80 W C KALB, C W KREIMENDAHL, D C BUSBY N00014-76-C-0390

UNCLASSIFIED TR-107

NL

[of]
ALL
AGENCY



END
DATE
FILMED
4-80
DDP

(14) TR-107

OFFICE OF NAVAL RESEARCH

(15) Contract No. N00014-76-C-0390

Task No. NR 053-608

(9) TECHNICAL REPORT NO. 107

LEVEL IV

DTIC
ELECTE
MAR 3 1980

ADA081321

(6) B(11)
A NMR Study of the Closo-hydridorhodacarborane $3,3-[(C_2H_5)_3P]_2-3-H-3,1,2-$
 $RhC_2B_9H_{11}(Ia)$ Prepared in situ from $3,3-(Ph_3P)_2-3-H-3,1,2-RhC_2B_9H_{11}(I)$.

by

(10) W. C. Kalb, C. W. Kreimendahl, D. C. Busby, M. F. Hawthorne

Prepared for Publication
in
Inorganic Chemistry

DTIC
ELECTE
MAR 3 1980

DDC FILE COPY

(12) 21

Department of Chemistry
University of California
Los Angeles, California 90024

(11) 22 Jan 1980

Reproduction in whole or part is permitted
for any purpose of the United States Government

Approved for Public Release; Distribution Unlimited

072255. AM
80 2 28 024

A ^{11}B NMR Study of the Closo-hydridorhodacarborane 3,3- $[(\text{C}_2\text{H}_5)_3\text{P}]_2\text{-3-H-3,1,2-}$
 $\text{RhC}_2\text{B}_9\text{H}_{11}(\text{Ia})$ Prepared in situ from 3,3- $(\text{Ph}_3\text{P})_2\text{-3-H-3,1,2-RhC}_2\text{B}_9\text{H}_{11}(\text{I})$

by

W. C. Kalb, C. W. Kreimendahl, D. C. Busby, and M. F. Hawthorne*

Department of Chemistry
University of California
Los Angeles, California 90024

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

SUMMARY

↙
A complete assignment of the ^{11}B NMR spectrum (111.8 MHz) of 3,3- $(\text{C}_2\text{H}_5)_3\text{P}$ $_2$ -3-H-3,1,2- $\text{RhC}_2\text{B}_9\text{H}_{11}$ (Ia) is proposed on the basis of ^{11}B NMR data obtained from specifically deuterated derivatives of I, and from 3,3- $(\text{Ph}_3\text{P})_2$ -3-H-9,12-(Br) $_2$ -3,1,2- $\text{RhC}_2\text{B}_9\text{H}_9$ and 6-Ph-3,3- $(\text{Ph}_3\text{P})_2$ -3-H-3,1,2- $\text{RhC}_2\text{B}_9\text{H}_{10}$. The ^{11}B NMR of ~~3,3- $(\text{C}_2\text{H}_5)_3\text{P}$ $_2$ -3-H-3,1,2- $\text{RhC}_2\text{B}_9\text{H}_{11}$~~ ^{compound 12 ABOVE} in tetrahydrofuran consists of five resonances of relative intensity 1:1:2:2:3 (reading upfield) which are assigned to B(10), B(8), B(9,12), B(4,7), B(6) and B(5,11), respectively.

Introduction

The first metallocarborane catalyst for the hydrogenation and isomerization of olefins, $3,3-(\text{Ph}_3\text{P})_2-3\text{-H-3,1,2-RhC}_2\text{B}_9\text{H}_{11}(\text{I})$,¹ also catalyzes the exchange of terminal boron-hydrogen bonds for deuterium in the presence of D_2 .^{2,3} These exchanges were found to proceed for a wide variety of carborane and metallocarborane substrates and, during these exchanges, deuteration also occurs at B-H vertices in I. In order to determine the specificity for self deuteration of I the ^{11}B NMR study presented herein was undertaken. Our goal was to completely assign the ^{11}B NMR spectrum of I and relate this information to a mechanism for the B-H/D exchange catalyzed by I and other metallocarboranes.

The solid state structure of I has been determined by a single crystal X-ray diffraction study,⁴ and found to be that of a regular icosahedron with the rhodium vertex symmetrically bonded to the pentagonal face of the dicarbollide ligand. For reasons discussed below, the ^{11}B NMR spectrum of I consists of broad, poorly resolved resonances.¹ With the discovery⁵ that substitution of triethylphosphine ligands for the triphenylphosphine ligands in I is accompanied by a marked improvement in the ^{11}B NMR spectrum of the derived complex $3,3-[(\text{C}_2\text{H}_5)_3\text{P}]_2-3\text{-H-3,1,2-RhC}_2\text{B}_9\text{H}_{11}(\text{Ia})$, it became feasible to attempt an assignment of the spectrum.

An assignment of the ^{11}B NMR spectrum of the $1,2\text{-C}_2\text{B}_9\text{H}_{12}^{1-6}$ ion has been reported by Hawthorne et al⁷ and others.⁸ The ^{11}B NMR spectrum of $(\text{C}_2\text{B}_9\text{H}_{11})_2\text{Co}^-$ has been assigned by Todd and co-workers.⁸ The present study provides a complete assignment of the ^{11}B NMR spectrum of Ia. Details of the mechanism for the B-H/D exchange catalyzed by I will be reported at a later date.

Results and Discussion

While ^{11}B NMR spectroscopy has found widespread application in the chemistry of boron hydrides, metalloboranes and metallocarboranes, the ^{11}B NMR spectra of metallocarboranes with bulky dissociable ligands have proven somewhat less amenable to analysis. Broad resonances are often observed for these complexes, especially for those which contain two metal-bound triphenylphosphine ligands. It was determined however that substitution of triethylphosphine for triphenylphosphine in I greatly improved the ^{11}B NMR spectrum of the derived complex. The 111.8 MHz ^{11}B NMR spectrum of I and 3,3- $[(\text{C}_2\text{H}_5)_3\text{P}]_2$ -3-H-3,1,2- $\text{RhC}_2\text{B}_9\text{H}_{11}$ (Ia) in tetrahydrofuran (THF) solution are presented in Figure 1 for comparison.

Figure 1

The numbering system employed throughout this paper is presented in the right side of Figure 1.⁶

This dramatic difference between the ^{11}B NMR spectra of I and Ia is most probably due to changes in the rate of molecular tumbling for these two species in solution. The longitudinal relaxation times (T_1) are relatively short for boron nuclei due to the quadrupole moment of both ^{10}B and ^{11}B .⁹ The presence of bulky triphenylphosphine ligands in I apparently increases the molecular correlation time. This should lead to effective quadrupolar relaxation and thus to broad ^{11}B NMR resonances for I. In contrast, the presence of the less bulky triethylphosphine ligands in complex Ia could afford an increase in the rate of tumbling (corresponding to decrease in the molecular correlation time). This would provide an increase in T_1 , resulting in diminished ^{11}B NMR linewidths.^{10,11}

The ^{11}B NMR spectrum of Ia in THF solution indicates the presence of a mirror plane (the spectrum contains 5 resonances of relative area 1:1:2:2:3


reading upfield), this is indicative of a higher degree of symmetry for I in solution than is observed for I in the solid state.⁴ This increase in symmetry is accounted for by a rapid rotation of the $\{\text{Rh}(\text{L})_2(\text{H})\}$ vertex in Ia about the center of the pentagonal face of the dicarbollide ligand.¹²

The mirror symmetry indicated for Ia would lead ideally to the observation of six resonances in the ^{11}B NMR spectrum of relative intensity 1:1:1:2:2:2. The presence of the weight three resonance observed for Ia is attributed to coincidental overlap of the resonance for a unique boron (B(6), B(8), or B(10)) with the resonance due to one of the symmetry equivalent pairs (B(4,7), B(5,11), or B(9,12)) (*vide infra*).

It was determined during the course of this study that samples of I in THF solution, when treated with excess triethylphosphine, gave ^{11}B NMR spectra identical to those which are observed for complex Ia in the same solvent. The hydridorhodacarboranes in this study were prepared and characterized as the bis-tri-phenylphosphine complexes. The ^{11}B NMR spectra reported here are for the bis-triethylphosphine complexes prepared *in situ*.


4,5,6,7,11- D_5 -1,2- $\text{C}_2\text{B}_9\text{H}_7^{1-}$ was prepared by base degradation of 3,4,5,6,7,11- D_6 -1,2- $\text{C}_2\text{B}_{10}\text{H}_6$ according to a literature procedure.¹³ 6-Ph- $\text{C}_2\text{B}_9\text{H}_{11}^{1-}$ was prepared analogously from 6-Ph- $\text{C}_2\text{B}_{10}\text{H}_{11}$.¹⁴ 9,12-(Br) $_2$ -1,2- $\text{C}_2\text{B}_9\text{H}_{10}^{1-}$ was prepared by base degradation of 9,12(Br) $_2$ -1,2- $\text{C}_2\text{B}_{10}\text{H}_{10}$.¹⁵ All monoanions were isolated as the tetramethylammonium salts and were dried over P_2O_5 *in vacuo*. μ -4,7,8,9,12- D_6 - $\text{C}_2\text{B}_9\text{H}_6^{1-}$ was prepared from K^+ 1,2- $\text{C}_2\text{B}_9\text{H}_{12}^{1-}$ in 6M DCl/D $_2\text{O}$ (6 days, 25°C).⁷ The ^{11}B NMR spectrum of this species revealed partial exchange had also occurred at B(5,11); only B(10), and B(6) retained complete coupling to terminal hydrogen, indicating these sites had not undergone exchange.

The derivatives of I were prepared from the appropriate 1,2- $\text{C}_2\text{B}_9\text{H}_{12}^{1-}$ derivative and $(\text{Ph}_3\text{P})_3\text{RhCl}$ in refluxing ethanol according to a procedure developed in this laboratory¹ (Equation 1). Typically, the reactions were complete after



10-15 minutes, at which time the hydridorhodacarboranes were isolated by Schlenk filtration, washed with ethanol and dried in vacuo.

No detectable scrambling of the deuterium label was observed in these experiments. This is due to both the short reaction times and the presence of triphenylphosphine (a product of reaction I) which has been shown to effectively inhibit the B-H/D exchange catalyzed by I.^{2,3} It is of note that I prepared from bridge deuterated 1,2-C₂B₉H₁₁D¹⁻ was found to contain only Rh-D ($\nu_{\text{Rh-D}}$ at 1520 cm⁻¹) with no trace of Rh-H detectable by IR (Equation 2). This conclusively establishes that the hydride in I results from the oxidative addition of



the bridging hydrogen in 1,2-C₂B₉H₁₂¹⁻.

The 111.8 MHz ¹¹B NMR spectrum of 4,5,6,7-11-D₅(Ia) in tetrahydrofuran, prepared in situ from 4,5,6,7,11-D₅(I) is shown in Figure 2a. Only the resonances at -2.8 ppm, -3.9 ppm and -9.0 ppm exhibit coupling to terminal hydrogens. The resonance of weight two at -9.0 ppm must therefore be due to the symmetry equivalent pair of boron atoms B(9,12). The resonance at -2.8 ppm and -3.9 ppm must be due to B(8), B(10) (indefinite order). The lack of B-H coupling in the weight three resonance at -21.0 ppm indicates this must contain the resonance for the remaining unique boron atom B(6).

This assignment is supported by the ¹¹B NMR spectrum of 6-Ph(Ia) prepared in situ from 6-Ph(I) in tetrahydrofuran solution (Figure 2b). Replacement of a phenyl group for hydrogen on boron causes the area 3 resonance at -20.3 ppm to decrease to an area of 2. The resonance due to B(6) is shifted downfield and appears as a singlet at -12.7 ppm.

The assignment of B(9,12) is supported by the ^{11}B NMR spectrum in THF of $9,12(\text{Br})_2(\text{Ia})$ prepared in situ from $9,12(\text{Br})_2(\text{I})$ (Figure 2c). The borons due to B(9,12) appear as a singlet at -3.2 ppm. Substitution of Br for terminal hydrogen on boron is generally accompanied by a shift to lower field of the ^{11}B NMR resonance due to the substituted boron.¹⁶ The resonances which remain unassigned in the spectrum of $9,12(\text{Br})_2(\text{Ia})$ exhibit chemical shifts nearly identical to those observed for Ia suggesting that the resonance at -9.0 ppm in the spectrum of Ia has moved to lower field in the spectrum of $(9,12)(\text{Br})_2(\text{Ia})$.

The ^{11}B NMR spectrum of Ia in THF prepared in situ from $3,4,7\text{-D}_3(\text{I})$ is shown in Figure 2d. The resonance of area two at -11.0 ppm appears as a singlet and is thus assigned to B(4,7). The resonance due to the remaining symmetry equivalent pairs of borons, B(5,11) must therefore occur at -21.1 ppm.

Only the resonances due to B(8), B(10) (which occur at low field) remain to be assigned. The ^{11}B NMR spectrum of $6,10\text{-H}_2(\text{Ia})$ in THF prepared in situ from $6,10\text{-H}_2(\text{I})$ is shown in Figure 2e. Only the resonance at -3.9 retains coupling to terminal hydrogen. The resonance due to B(6) has previously been shown to occur at -21.0 ppm (vide supra) and therefore the resonance at -3.9 ppm is assigned to B(10). The remaining resonance at -2.8 ppm is assigned to B(8) by elimination.

It is of interest to compare the present assignment with that of the analogous metallocarborane $(\text{C}_2\text{B}_9\text{H}_{11})_2\text{Co}^-$.⁸ The assignments (reading upfield) are for Ia: B(10), B(8), B(9,12), B(4,7), B(6), B(5,11); and for $(\text{C}_2\text{B}_9\text{H}_{11})_2\text{Co}^-$: B(8), B(10), B(9,12), B(4,7), B(5,11), B(6). All resonances in the ^{11}B NMR of Ia are found at higher field than those observed for $(\text{C}_2\text{B}_9\text{H}_{11})_2\text{Co}^-$ with the exception of B(6). Overall, however, the two spectra appear quite similar.

The different assignment for B(8) and B(10) for two such similar spectra should serve as a caution against assigning ^{11}B NMR resonances by comparison to "similar" species.


Experimental

Instrumentation

Infrared spectra were recorded on a Perkin-Elmer 137 Spectrometer. ^{11}B NMR (111.8 MHz) were obtained in the Fourier-transform mode on an instrument designed and constructed by Professor F. A. L. Anet of this department.

Materials

All solvents were reagent grade and, except where otherwise noted, were distilled under argon from the appropriate drying agent immediately prior to use. Ethanol (abs.) was saturated with nitrogen but otherwise used without further purification. Tetrahydrofuran and diethyl ether (Mallinckrodt) were dried over alumina and distilled from potassium. $(\text{Ph}_3\text{P})_3\text{RhCl}$ was prepared from $\text{RhCl}_3 \cdot 3\text{H}_2\text{O}$ (Matthey-Bishop) according to a procedure in the literature.¹⁷ $\text{K}^+7,8\text{-C}_2\text{B}_9\text{H}_{12}^-$ was prepared according to the literature¹⁸ and stored in a nitrogen-filled glove-box. DCl (20% in D_2O , 99 + atom % D, Alfa-Ventron) was diluted immediately prior to use with D_2O in a nitrogen-filled glove-bag.



$\text{K}^+1,2\text{-C}_2\text{B}_9\text{H}_{12}^-$ (1.0 g, 5.8 mmol) was weighed in the dry box into a 100 ml, round bottom flask equipped with a sidearm and stopcock. The flask was sealed with a gas inlet tube equipped with a stopcock, transferred to the bench, and fitted to a nitrogen line. D_2O (1 M in DCl , 15 ml) was syringed into the flask and the flask was shaken to achieve a homogeneous solution. The flask was then

sealed under a positive pressure of N_2 . After three days a sample of the solution was obtained with a syringe and transferred to a nitrogen-filled NMR tube. The ^{11}B NMR of this sample revealed complete deuteration at B(4,7) and the B-H-B bridge.

An additional 20 ml of D_2O was added to the flask, followed by $(CH_3)_4NCI$ (0.73 g, 6.7 mmol) in 5 ml of D_2O . The precipitate which formed was isolated by Schlenk-filtration, washed with three-10 ml portions of D_2O and dried at $80^\circ C$ over P_2O_5 in vacuo. A ^{11}B NMR of the product was obtained in CH_3CN .

Preparation of NMR samples of I and derivatives of I

Caution: $(C_2H_5)_3P$ is toxic: gloves should be worn and these operations should be performed in an efficient fume hood.

The sample of I (15 to 20 mg) was added to a 5 mm NMR tube which was then fitted with a rubber septum and purged with N_2 utilizing two syringe needles. After 5 mins, tetrahydrofuran (0.8 ml) was added to the sample with a syringe. Immediately thereafter, $(C_2H_5)_3P$ (0.15 ml) was added with a syringe. The needles were removed and the sample was shaken to dissolve the metallocarborane. ^{11}B NMR spectra of the samples so prepared were recorded after a period of 45 mins, to allow for complete phosphine exchange.

Acknowledgement

This research was supported in part by the National Science Foundation (Grant No. CHE77-22910) and the Office of Naval Research. We thank Professor F. A. L. Anet for the use of his NMR facility, and Dr. Dan Donovan for his technical assistance.

References and Notes

1. T. E. Paxson and M. F. Hawthorne, J. Am. Chem. Soc., **96**, 4674 (1974).
2. E. L. Hoel and M. F. Hawthorne, J. Am. Chem. Soc., **96**, 4676 (1974).
3. E. L. Hoel, M. Talebinasab-Savari and M. F. Hawthorne, J. Am. Chem. Soc., **99**, 4356 (1977).
4. G. E. Hardy, K. P. Callahan, C. E. Strouse, and M. F. Hawthorne, Acta Cryst. (1976), **B32**, 264.
5. The complex $3,3-[(C_2H_5)_3P]_2-3-H-3,1,2-RhC_2B_9H_{11}$ was first synthesized and completely characterized by R. T. Baker of these laboratories.
6. For continuity, an older numbering system is used to describe the position of substitution in $7,8-C_2B_9H_{12}^-$ (referred to as $1,2-C_2B_9H_{12}^-$ in this paper). Under this system, the numerical prefix which assigns the position of substitution, is the same for both the $C_2B_9H_{12}^-$ derivatives and the respective derivatives of I.
7. D. V. Howe, C. J. Jones, R. J. Wiersema, and M. F. Hawthorne, Inorg. Chem., **10**, 2516 (1971).
8. A. R. Siedle, G. M. Bodner, and L. J. Todd, J. Organomet. Chem., **33**, 137 (1971).
9. S. G. Shore, in "Boron Hydride Chemistry" (E. L. Muetterties, Ed.), p. 95, Academic Press, New York, 1975.
10. E. D. Becker, "High Resolution NMR", p. 206, Academic Press, New York, 1969.
11. This explanation has been offered to account for the broad ^{11}B NMR resonances found in metalloboranes which contain bulky aryl phosphines. J. D. Kennedy and J. Staves, Z. Naturforsch., **34b**, 808 (1979); and E. L. Muetterties, Rev. Pure Appl. Chem., **29**, 585 (1972).
12. T. B. Marder, R. T. Baker, and J. A. Doi, to be published.
13. We are grateful to Dr. E. L. Hoel for the sample of $3,4,5,6,7,11-D_6-1,2-C_2B_{10}H_6$ which was prepared from $1,4-5,6,7,8,9,10-D_{10}-B_{10}H_4$ (J. A. Dupont and M. F. Hawthorne, J. Am. Chem. Soc., **84**, 1804 (1962)) according to the procedure of L. J. Todd, Inorg. Synth., **XI**, 19 (1968).
14. M. F. Hawthorne and P. A. Wegner, J. Am. Chem. Soc., **90**, 896 (1968).

15. H. D. Smith, T. Knowles, and H. A. Schroeder, Inorg. Chem., 4, 107 (1965).
16. G. R. Eaton and W. N. Lipscomb, "NMR Studies of Boron Hydrides and Related Compounds", p. 407-408, W. A. Benjamin, Inc., New York, 1969.
17. J. A. Osborn, F. H. Jardine, J. F. Young, and G. Wilkinson, J. Chem. Soc. A, 1711 (1966).
18. M. F. Hawthorne, D. C. Young, P. M. Garrett, D. A. Owen, S. G. Schwerin, F. N. Tebbe, and P. A. Wegner, J. Am. Chem. Soc., 90, 862 (1968).

Figure 1 111.8 MHz ^{11}B NMR spectra of 3,3- L_2 -3-H-3,1,2- $\text{RhC}_2\text{B}_9\text{H}_{11}$

a) $\text{L} = \text{PPh}_3$ (I)

b) $\text{L} = \text{P}(\text{C}_2\text{H}_5)_3$ (Ia)












Figure 2

111.8 MHz ^{11}B NMR spectra of:

- a) 4,5,6,7,11- D_5 -(Ia)
- b) 6-Ph-(Ia)
- c) 9,12- Br_2 -(Ia)
- d) 3,4,7- D_3 -(Ia)
- e) 6,10- H_2 -(Ia)



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report 107	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) "A ^{11}B Study of the Closo-hydridorhodacarborane 3,3- [(C ₂ H ₅) ₃ P] ₂ -3-H-3,1,2-RhC ₂ B ₉ H ₁₁ (Ia) Prepared <u>in situ</u> from 3,3-(Ph ₃ P) ₂ -3-Y-3,1,2-RhC ₂ B ₉ H ₁₁ (I)"		5. TYPE OF REPORT & PERIOD COVERED Interim
7. AUTHOR(s) W. C. Kalb, C. W. Kreimendahl, D. C. Busby, and M. F. Hawthorne		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS The University of California Department of Chemistry Los Angeles, California 90024		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0390
11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Branch Office of Naval Research Washington, D.C. 20360		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-608
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE January 22, 1980
		13. NUMBER OF PAGES 12 (incl. figs.)
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Carborane Nuclear Magnetic Resonance (NMR) Metallocarborane Rhodacarborane Boron		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A complete assignment of the ^{11}B NMR spectrum (111.8 MHz) of 3,3-[(C ₂ H ₅) ₃ P] ₂ -3-H-3,1,2-RhC ₂ B ₉ H ₁₁ (Ia) is proposed on the basis of ^{11}B NMR data obtained from specifically deuterated derivatives of I, and from 3,3-(Ph ₃ P) ₂ -3-H-9,12-(Br) ₂ -3,1,2-RhC ₂ B ₉ H ₉ and 6-Ph-3,3-(Ph ₃ P) ₂ -3-H-3,1,2-RhC ₂ B ₉ H ₁₀ . The ^{11}B NMR of 3,3-[(C ₂ H ₅) ₃ P] ₂ -3-H-3,1,2-RhC ₂ B ₉ H ₁₁ in tetrahydrofuran consists of five resonances of relative intensity 1:1:2:2:3 (reading upfield) which are assigned to B(10), B(8), B(9,12), B(4,7), B(6) and B(5,11), respectively.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

APPENDIX

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Branch Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Branch Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. R. N. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1	Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1
Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1	Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1
		Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1	Professor O. T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214	1
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1	Professor P. S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1	Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1	Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
Professor H. Abrahamson University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1	Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1